

RESEARCH

Open Access

Pathological alpha-synuclein propagates through neural networks

Masami Masuda-Suzukake¹, Takashi Nonaka¹, Masato Hosokawa², Maki Kubo¹, Aki Shimozawa¹, Haruhiko Akiyama² and Masato Hasegawa^{1*}

Abstract

Background: α -Synuclein is the major component of filamentous inclusions that constitute the defining characteristic of Parkinson's disease, dementia with Lewy bodies and multiple system atrophy, so-called α -synucleinopathies. Recent studies revealed that intracerebral injection of recombinant α -synuclein fibrils into wild-type mouse brains induced prion-like propagation of hyperphosphorylated α -synuclein pathology. However, the propagation mechanisms of α -synuclein have not been fully elucidated.

Results: In this study, in order to establish where and how α -synuclein pathology propagates, we injected recombinant mouse α -synuclein fibrils into three different brain areas (substantia nigra, striatum, and entorhinal cortex) of wild-type mice and compared the resulting distributions of α -synuclein pathology at 1 month after injection. Distinct patterns of pathology were observed in mice injected at the different sites. Within one month after injection, the pathology had spread to neurons in areas far from the injection sites, especially areas with direct neural connections to the injection sites. Surprisingly, phosphorylated tau and TDP-43 pathologies were also observed in mice injected with α -synuclein fibrils into striatum and entorhinal cortex at one month after injection. Phosphorylated tau and TDP-43 were accumulated in dot-like inclusions, but these were rarely colocalized with α -synuclein pathology. It seems that accumulation of α -synuclein has a synergistic effect on tau and TDP-43 aggregation. Additionally, intracerebral injection with sarkosyl-insoluble fraction prepared from wild-type mice injected synthetic α -synuclein fibrils can also induce phosphorylated α -synuclein pathology in wild-type mice.

Conclusions: Our data indicate that α -synuclein aggregation spread by prion-like mechanisms through neural networks in mouse brains.

Keywords: α -Synuclein, Lewy bodies, Propagation, Prion

Introduction

Parkinson's disease (PD) and dementia with Lewy bodies (DLB) are progressive neurodegenerative diseases characterized by appearance of Lewy bodies (LBs) and Lewy neurites (LNs) [1]. α -Synuclein (α syn) is the major component of LBs and LNs, and is deposited in a hyperphosphorylated form in β -sheet-rich amyloid fibrils [2-5]. Five missense mutations in the α syn gene and occurrence of gene multiplication have been identified in the familial forms of PD and DLB [6-13]. Moreover, it was reported that the mutations affect amyloid fibril formation *in vitro*,

either accelerating fibril formation [14-16] or resulting in formation of fibrils that are more fragile and easy to propagate than wild-type (WT) fibrils [17]. These results clearly indicate that abnormalities of α syn can induce PD and DLB. Distribution of α syn pathology in brains with sporadic PD occurs from olfactory bulb and/or brainstem, and spreads to other brain regions concomitantly with progression of disease symptoms [18,19]. Thus, spread of α syn pathology in the brain can be regarded as the underlying mechanism of progression of these diseases. Recently intracerebral injection of synthetic α syn fibrils and/or insoluble α syn from diseased brain was shown to induce α syn pathology that propagated throughout the brain in a prion-like manner in WT mouse [20,21], α syn transgenic mouse [22-24] and monkey [25]. However, the mechanisms through which exogenous abnormal fibrils enter

* Correspondence: hasegawa-ms@igakuken.or.jp

¹Department of Neuropathology and Cell Biology, Tokyo Metropolitan Institute of Medical Science, 2-1-6 Kamikitazawa, Setagaya-ku, Tokyo 156-0057, Japan

Full list of author information is available at the end of the article

neurons and through which insoluble α syn is transported to other neurons remain unknown.

To investigate where α syn pathologies develop and how they propagate, we injected recombinant α syn fibrils into substantia nigra, striatum, or entorhinal cortex of WT mice, and compared the spreading patterns and distribution of phosphorylated α syn pathologies at 1 month after intracerebral injections. Our results clearly suggest that propagation of pathological α syn occurred along neural circuits and involved trans-synaptic transport. We also showed that α syn pathology induced tau and TDP-43 accumulation in WT mice, similar to that seen in DLB brains. This mouse model should be useful for elucidating mechanisms of disease progression of synucleinopathy and also for development of novel disease-modifying drugs.

Materials and methods

Antibodies

Antibodies used in this study are summarized in Additional file 1: Table S1. 1175 polyclonal antibody was raised against an α syn peptide phosphorylated at serine 129 [21,26]. Anti-phosphorylated α syn mouse monoclonal antibody, #64 [5] and anti-human α syn specific mouse monoclonal antibody, LB509 [27] were kindly provided from Dr. Iwatsubo.

Rabbit polyclonal pS396 antibody (Calbiochem) is specific for phosphorylated tau at serine 396; biotin-AT8 (Thermo Scientific) is specific for phosphorylated tau at serine 202/threonine 205; anti-mouse α syn rabbit monoclonal antibody (Cell Signaling Technology) is specific for mouse α syn. Rabbit polyclonal pTDP-43 antibody is specific for phosphorylated at serine 409/410 [28].

Preparation of recombinant α syn monomer and fibrils

Mouse α syn cDNA in bacterial expression plasmid pRK172 was used. α Syn were expressed in *Escherichia coli* BL21 (DE3) cells and purified using boiling, Q-sepharose ion exchange chromatography and ammonium sulfate precipitation. Purified α syn protein was dialyzed against 30 mM Tris-HCl, pH 7.5, and cleared using ultracentrifugation at 113,000 g for 20 min. Protein concentration was determined by reverse phase HPLC. Proteins were loaded on an Aquapore RP-300 column (PerkinElmer Brownlee) equilibrated in 0.09% trifluoroacetic acid with linear gradient of acetonitrile 0 to 50% at a flow rate of 1 ml/min [21]. Purified mouse α syn monomer (7 mg/ml) in 30 mM Tris-HCl, pH 7.5, containing 0.1% NaN_3 was incubated at 37°C in a shaking incubator at 200 rpm for 72 h. α Syn fibrils were pelleted by spinning at 113,000 g for 20 min and




Figure 1 Distribution of phosphorylated α syn pathology in α syn fibril-injected mice at 1 month after injection. **(A)** Injection into SN induced α syn pathology mainly in SN (3.08 mm posterior to bregma), amygdala (1.58 mm posterior to bregma) and stria terminalis (0.02 mm anterior to bregma). **(B)** Injection into Str induced severe α syn pathology throughout the brain, including Str (0.26 mm anterior to bregma), amygdala (1.58 mm posterior to bregma), SN (2.70 mm posterior to bregma) and a wide range of cortex. **(C)** Injection into EC induced α syn pathology that was concentrated in EC (3.52 mm posterior to bregma), dentate gyrus (3.52 mm posterior to bregma), CA3 (3.52 mm posterior to bregma), fimbria (1.94 mm posterior to bregma), and septal nucleus (0.02 mm anterior to bregma). Blue-dashed box and red dots indicate the injection site and psyn pathology, respectively.

suspended in PBS. α Syn fibrils were sonicated with an ultrasonic homogenizer (VP-5S, TAITEC) before use. To determine the concentration, fibrils were dissolved in 8 M guanidine hydrochloride and analyzed by RP-HPLC as described above.

Mice

C57BL/6 J mice, used as WT mice, were purchased from CLEA Japan, Inc. α Syn (SNCA) knockout mice [29] were purchased from the Jackson Laboratory.

Stereotaxic surgery

Four- to six-month-old mice anesthetized with 50 mg/kg pentobarbital sodium were unilaterally injected with 10 μ g of recombinant mouse α syn fibrils into substantia nigra (SN, $n = 6$) (A-P: -3.0 mm; M-L: -1.3 mm; D-V: -4.7 mm from the bregma and dura) [21], striatum (Str, $n = 6$) (A-P: 0.2 mm; M-L: -2.0 mm; D-V: -2.6 mm) [20], or entorhinal cortex (EC, $n = 6$) (A-P: -3.1 mm; M-L: -4.0 mm; D-V: -2.7 mm). Mice were anesthetized with isoflurane and killed by decapitation. For immunohistochemistry (IHC, $n = 3$), brains were fixed in 10% formalin neutral buffer solution (Wako). For biochemical analysis ($n = 3$), brains were snap-frozen on dry ice and stored at -80°C . All experimental protocols were approved by the Animal

Care and Use Committee of the Tokyo Metropolitan Institute of Medical Science.

Peripheral injection of asyn

For intraperitoneal injection, 2-month-old C57BL/6 J mice were injected intraperitoneally with 100 μ g of mouse α syn monomer or fibrils. At 6 months after injection, the pathology of mouse brains in both groups ($n = 3$ each) was tested by immunohistochemistry (IHC). For oral administration, 2- or 3-month-old C57BL/6 J mice were orally administered with 400 μ g of human α syn monomer, human α syn fibrils, mouse α syn monomer or mouse α syn fibrils every two weeks for 4 times. At 12 months post final administration, pathology in mouse brains ($n = 3$ each) was analyzed by IHC.

Immunohistochemistry

Fixed brains were cut on a vibratome (Leica) at 50 μ m thickness. For high-sensitivity detection, mouse brain sections were treated with formic acid for 30 min, washed, and boiled at 100°C for 30 min. The sections were then incubated with 0.5% H_2O_2 in methanol to inactivate endogenous peroxidases, blocked with 10% calf serum in PBS, and immunostained with appropriate antibodies. After incubation with the biotinylated-secondary antibody (Vector), labeling was detected using the ABC staining kit (Vector).




Figure 2 Staining of WT mouse brains injected with α syn fibrils at 1 month after injection by using 1175 antibody. (A) Psyn pathology in mice injected into SN. (B) Psyn pathology in mice injected into Str. (C) Psyn pathology in mice injected into EC. SN: substantia nigra, Amy: amygdala, ST: stria terminalis, Str: striatum, Ctx: cortex, EC: entorhinal cortex, DG: dentate gyrus. Scale bar represents 50 μ m.

Confocal microscopy

For double-label immunofluorescence to detect phosphorylated α syn and tau, brain sections were incubated overnight at 4°C in a cocktail of #64 antibody and anti-pS396 antibody. The sections were washed and incubated in a cocktail of Alexa568-conjugated goat anti mouse IgG (Molecular Probes) and Alexa488-conjugated goat anti rabbit IgG (Molecular Probes). After further washing, sections were stained with TOPRO-3, cover-slipped with Vectashield (Vector) and observed with a

laser-scanning confocal fluorescence microscope (LSM5 PASCAL; Carl Zeiss).

Biochemical analysis

Biochemical analysis of mouse brains ($n = 3$ per group) was conducted as described previously [21]. Briefly, brains were homogenized in 20 volumes (w/v) of buffer A (10 mM Tris-HCl, pH 7.4, 0.8 M NaCl, 1 mM EGTA and 10% sucrose), then spun at 100,000 g for 30 min at 4°C, and the supernatant was retained as buffer-soluble




Figure 3 Induction of phosphorylated tau inclusions in WT mice injected with α syn fibrils at 1 month after injection. (A) Phosphorylated tau was not accumulated in mice injected into SN, based on staining with anti-pS396 antibody. (B-C) Dot-like tau inclusions were observed in mice injected into Str by using anti-pS396 antibody (B) and AT8 (C) (D-E). Dot-like tau inclusions were also detected in mice injected into EC by using anti-pS396 antibody (D) and AT8 (E). SN: substantia nigra, Amy: amygdala, ST: stria terminalis, Str: striatum, Ctx: cortex, EC: entorhinal cortex, DG: dentate gyrus. Scale bar represents 50 μ m.

fraction. The pellet was homogenized in 20 volumes of buffer A containing 1% Triton X-100 and incubated for 30 min at 37°C. After centrifugation at 100,000 g, the Triton-insoluble pellet was further homogenized in buffer A containing 1% sarkosyl and incubated at 37°C for 30 min. Samples were spun at 100,000 g for 30 min. The sarkosyl-pellet was sonicated in 30 mM Tris-HCl, pH 7.4, and used for immunoblotting as sarkosyl-insoluble fraction. The samples were subjected to SDS-PAGE and proteins were electrotransferred onto a polyvinylidene difluoride membrane, probed with appropriate antibodies and detected as described previously [21].

Behavioral tests

For behavioral tests, C57BL/6 J male mice were used. Mouse α syn fibrils (10 μ g) were injected into SN (n = 10), Str (n = 15), or EC (n = 14) of 3-month-old mice, and the same amount of mouse α syn monomer was injected into SN (n = 9), Str (n = 8), or EC (n = 8) of control mice. At 3 months after injection, motor and cognitive activities were evaluated as described below.

Rotarod test

Motor coordination and balance were measured in terms of performance on the rotarod. Mice were placed on 3-cm

diameter rods and the speed of the rotation was increased from 0 to 40 rpm over 5 min. Latency to fall was recorded. Each mouse was tested three times and the average was used. Statistical analyses were performed using Student's *t*-test.

Wire hang test

Neuromuscular abnormalities were tested with the wire hang test. The mouse was placed on a wire cage lid, which was waved gently so that the mouse gripped the wire and then inverted. Latency to fall was recorded with a 300-sec cut-off time. The test was conducted three times and statistical analyses were performed using Student's *t*-test.

Y-maze test

The Y-maze apparatus (Muromachi kikai) consisted of three arms (40 cm \times 3 cm) made of grey plastic joined in the middle to form a Y shape. Mice were placed into one of the arms of the maze and allowed to freely explore the three arms for an 8-min session. Alternation between arms was recorded. The Y-maze test was conducted twice. Statistical analyses were performed using Student's *t*-test.




Figure 4 Induction of phosphorylated TDP-43-positive structures in WT mice injected with α syn fibrils at 1 month after injection. (A) Phosphorylated TDP-43 was not accumulated in mice injected into SN, based on staining with anti-pS409/410 antibody. (B) Dot-like TDP-43 inclusions were observed in mice injected into Str. (C) Dot-like TDP-43 inclusions were also detected in mice injected into EC. SN: substantia nigra, Amy: amygdala, ST: stria terminalis, Str: striatum, Ctx: cortex, EC: entorhinal cortex, DG: dentate gyrus. Scale bar represents 50 μ m.

Transmission experiments

Recombinant human α syn fibrils (10 μ g) were injected into SN of 4-month-old WT mice ($n = 4$). At 9 months after injection, sarkosyl-insoluble pellets were prepared from the whole brains as described above, collected in one tube, and stored at -80°C until use. Sarkosyl-insoluble pellets were suspended in 100 μ l PBS and sonicated for 30 seconds (TAITEC, VP-5S), and 5- μ l aliquots were injected into Str of 4-month-old WT mice ($n = 10$). At 3 months post injection, pathology was analyzed by IHC.

Results

We investigated the spread of α syn pathology in brains of mice after unilateral injection of recombinant mouse α syn fibrils into SN, Str, or EC. We confirmed the purity of recombinant α syn monomer and fibrils used in this study; they didn't contain any contaminants (Additional file 1: Figure S1). Using highly sensitive immunohistochemistry (IHC) with anti-phosphorylated α syn (psyn) antibody 1175, we evaluated α syn pathology in the brains at 1 month after injection. The distribution of α syn pathology observed in these mice is illustrated in Figure 1. In mice injected into SN (Figure 1A), abnormal psyn pathology was restricted mainly to SN (3.08 mm posterior to bregma), amygdala (1.58 mm posterior to bregma), and stria terminalis (0.02 mm anterior to bregma) of the

hemisphere on the injection side. In these mice, psyn was accumulated in neurites and soma (Figure 2A). In mice injected into Str, psyn pathology was widely distributed bilaterally throughout the brain, including Str (0.26 mm anterior to bregma), amygdala (1.58 mm posterior to bregma), SN (2.70 mm posterior to bregma) and cortex (Figure 1B). Psyn pathology was accumulated mainly in neurites, and partly in soma (Figure 2B). Injection of α syn fibrils into EC induced severe psyn pathology in EC (3.52 mm posterior to bregma), dentate gyrus (3.52 mm posterior to bregma), hippocampal CA3 region (1.94 and 3.52 mm posterior to bregma), fimbria (1.94 mm posterior to bregma), and septal nuclei (0.02 mm anterior to bregma) on the injection side, as well as moderate psyn pathology in hippocampus on the contralateral side (Figure 1C). Psyn pathology was mainly observed in neurites and perinuclear regions (Figure 2C). No such psyn accumulation was detected in α syn KO mice injected with α syn fibrils into Str (Additional file 1: Figure S2A). Thus, there are major differences among these mice in the development and spread of α syn pathologies, demonstrating that the propagation pattern depends upon the injection site.

To investigate whether other pathologies are also induced by the injection of α syn fibrils, we performed IHC analysis using anti-tau, anti-TDP and anti-A β antibodies. No tau pathology was observed in mice injected into SN




Figure 5 Dot-like ptau-positive structures showed little colocalization with psyn pathology in hippocampus of mice injected with α syn fibrils into EC at 1 month after injection. Brain sections were double-labeled with anti-psyn antibody (#64, red) and anti-ptau antibody (pS396, green). Scale bar represents 50 μ m.





Figure 7 Fibril-injected mice showed motor dysfunctions compared to monomer-injected mice at 3 months after injection. (A) Rotarod test. Mice injected into SN and Str showed lower performance in the rotarod test. **(B)** Wire hang test. **(C)** Y-maze test. All error bars indicates mean \pm S. E. M. * $p < 0.01$, ** $p < 0.05$. M: asyn monomer-injected, F: asyn fibril-injected.

Discussion

Luk *et al.* and we have established that insoluble α syn shows prion-like propagation behavior in WT mouse brain [20,21], but the mechanism of spreading remains poorly understood. In this study, we investigated the spread and distribution pattern of psyn pathology in mouse brain injected with recombinant α syn fibrils into three different brain regions: SN, Str, and EC. We assessed the distribution at 1 month post injection by using highly sensitive IHC. Pretreatment of brain sections with formic acid and heat enabled detection of psyn pathology at only 1 month after injection. When α syn fibrils were injected into SN, psyn pathology only appeared in the central nucleus of amygdala and stria terminalis, which are located far from SN, while there was no detectable psyn pathology around SN (Figure 1A). Amygdala is connected with SN [30], and stria terminalis serves as a major output pathway of the central nucleus of amygdala. These findings strongly indicate that spreading of psyn pathology does not occur by simple diffusion or nonspecific transport. In the case of injection into Str, psyn pathology was observed in amygdala, SN and a

wide range of cortices (Figure 1B). Str has direct projection to SN and amygdala [31], and many parts of the neocortex innervate the Str [32]. Injection into EC induced pathology in EC, dentate gyrus, hippocampal CA3, fimbria and septal nucleus (Figure 1C). Dentate gyrus receives projection from EC via the perforant pathway, and septal nucleus and fimbria have direct connections with hippocampus. Therefore, the data strongly suggest that propagation of pathological α syn occurs via axonal transport and a trans-synaptic pathway, in accordance with reports that α syn fibrils can be internalized by neurons and transferred from axons to second-order neurons in culture [33] and in animal models [34,35]. In patients with sporadic PD, the distribution of Lewy bodies and Lewy neurites seems to spread retrogradely [18,36]. In the present study, focusing on Str and SN, injection with α syn fibrils into Str induced α syn pathology in SN at 1 month after injection (Figure 1B). However, injection into SN did not induce pathology in Str at that time (Figure 1A), and pathology only became apparent in Str at 3 months after injection [21], indicating there is a dominant direction of spread. Propagation from EC to



dentate gyrus via the perforant pathway might occur via anterograde transport (Figure 1C). At least under our experimental conditions, propagation of α syn seems to occur via both anterograde and retrograde transport processes. Thus, the predominant direction of spread presumably depends on cell types or brain areas. Similarly, tau is

also reported to propagate via a trans-synaptic pathway in animal models [37,38]. Thus, axonal transport and trans-synaptic transport appear to be common pathways of propagation of intracellular aggregated proteins.

In addition, we found that tau and TDP-43 accumulation also occurred in WT mice injected with α syn fibrils

into Str and EC at 1 month after injection (Figures 3B-E and 4). The morphological patterns of tau and TDP-43 accumulation were apparently different from that of α syn pathology (Figures 2, 3 and 4) and there was little colocalization (Figure 5), as in DLB brains [39,40] and a DLB mouse model [41]. Recently, Guo et al. reported that there are two strains of recombinant α syn fibrils, strains A and B, and the two strains differently affect tau inclusion formation [42]. They reported that strain A (preformed fibrils) only infrequently induced tau inclusions and psyn pathology showed little colocalization with tau inclusions, whereas strain B (generated through repetitively seeded fibrillization *in vitro*) efficiently induced tau inclusions that were highly colocalized in neurons. The α syn fibrils we used in this study are similar to strain A, and in agreement with their work [42], we also detected small amounts of tau inclusions that showed little colocalization with α syn pathology. In addition, tau accumulation was not observed in α syn KO mice injected with α syn fibrils by biochemical analysis (Additional file 1: Figure S4) or IHC (Additional file 1: Figure S2B). Thus, tau accumulation was induced by α syn accumulation, and this occurred through a synergistic effect rather than a cross-seeding effect. The tau accumulation might be caused by a secondary effect of α syn accumulation, such as dysfunction of cellular activity [43,44] or abnormality in protein degradation machinery [45].

Biochemical analysis clearly showed that accumulated α syn was phosphorylated and ubiquitinated similarly to that in DLB brain, regardless of injection site (Figure 6). This indicates that injection with the same fibrils as seeds induces α syn aggregation in the same fashion, as is the case with prion strains.

Furthermore, α syn fibril-injected mice showed modest motor abnormalities compared to the monomer-injected mice at 3 months after injection (Figure 7). This strongly suggests that propagation of psyn pathology induced motor phenotypes, although we could not detect cognitive dysfunction in the Y-maze test. It is possible that this is because we injected α syn fibrils unilaterally, and the functions of the contralateral side of brain might be well maintained. In our previous study, we could not detect any abnormalities in fibril-injected mice at 6 months after injection [21]. The discrepancy may have arisen from differences in the test conditions, because in the previous study, we used female mice injected with human α syn fibrils, whereas in this study we used male mice injected with mouse α syn. Mouse α syn fibrils propagate more efficiently in WT mice than do human fibrils [21], and we think this was the main reason why we could detect motor abnormalities in the present study.

We next examined if insoluble α syn accumulated in WT mice shows transmissibility. Our results demonstrate that insoluble α syn accumulated in WT mice can induce α syn pathology in other WT mice (Figure 8C, D),

analogously to prion transmission. We also examined intraperitoneal injection or oral administration with α syn fibrils into WT mice (see Materials and methods), but failed to detect any psyn pathology in the central nervous system at 6 months or 14 months after injection, respectively (data not shown).

Conclusions

Intracerebral injection with α syn fibrils into WT mouse brains enables to induce phosphorylated α syn pathology and the distribution of pathology depends on the injection sites. Furthermore, α syn pathology has a synergistic effect on tau and TDP-43 aggregation. We conclude that α syn fibrils have prion-like transmissibility and it might spread via axonal and trans-synaptic transports in mouse brains.

Additional file

Additional file 1: Table S1. Antibodies used in this study. **Figure S1.** HPLC charts of recombinant mouse α syn used in this study. Mouse syn monomer showed only one peak that is derived from msyn monomer. Mouse syn fibril gave two peaks at 0.14 min (guanidine HCl) and at 6.8 min (msyn fibril). **Figure S2.** α Syn and tau accumulation was never observed in fibril-injected α syn KO mice at 3 months after injection. **(A)** No psyn-positive pathology was observed with 1175 antibody. **(B)** Tau accumulation was not detected with pS396 antibody. Str: striatum, Amy: amygdala, SN: substantia nigra, sensory ctx: sensory cortex. Scale represents 50 μ m. **Figure S3.** A β accumulation was not observed in α syn fibril-injected WT mice at 1 month post injection. Sections were stained with anti-mouse A β antibody. Mice injected into SN **(A)**, Str **(B)**, and EC **(C)**. SN: substantia nigra, Amy: amygdala, ST: stria terminalis, Str: striatum, Ctx: cortex, EC: entorhinal cortex, DG: dentate gyrus. Scale represents 50 μ m. **Figure S4.** Biochemical analysis of α syn KO mice injected with human α syn fibrils. The brain was divided into two parts at the longitudinal fissure of the cerebrum. Sarkosyl-insoluble fractions were obtained from the right and left brains, and analyzed by immunoblotting with #64, LB509 or anti-mouse α syn antibodies. Exogenous human α syn fibrils were detected in sarkosyl-insoluble fractions and were not phosphorylated at 0 and 7 days after injection. They were subsequently degraded and disappeared within 30 days post injection. Phosphorylated α syn accumulation was never observed at 90 days after injection.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

MMS performed stereotaxic surgery, biochemical and IHC studies and wrote the manuscript. MK and AS performed IHC analysis and behavioral tests. TN and MH helped for interpretation of data. HA provided antibodies and helpful advice. MH performed study design and wrote the manuscript. All authors read and approved the final manuscript.

Acknowledgements

This work was supported by MEXT KAKENHI Grant Numbers 12937622, 12901980 (to M.H.), JSPS KAKENHI Grant Number 23700433 (to M.M.-S.) and MHLW Grant Number 12946221 (to M.H.).

Author details

¹Department of Neuropathology and Cell Biology, Tokyo Metropolitan Institute of Medical Science, 2-1-6 Kamikitazawa, Setagaya-ku, Tokyo 156-0057, Japan. ²Dementia Research Project, Tokyo Metropolitan Institute of Medical Science, Setagaya-ku, Tokyo, Japan.

Received: 14 July 2014 Accepted: 18 July 2014

Published: 6 August 2014

References

- Goedert M (2001) Alpha-synuclein and neurodegenerative diseases. *Nat Rev Neurosci* 2(7):492–501, doi:10.1038/35081564
- Spillantini MG, Schmidt ML, Lee VM, Trojanowski JQ, Jakes R, Goedert M (1997) Alpha-synuclein in Lewy bodies. *Nature* 388(6645):839–840, doi:10.1038/42166
- Spillantini MG, Crowther RA, Jakes R, Hasegawa M, Goedert M (1998) Alpha-Synuclein in filamentous inclusions of Lewy bodies from Parkinson's disease and dementia with lewy bodies. *Proc Natl Acad Sci U S A* 95(11):6469–6473
- Baba M, Nakajo S, Tu PH, Tomita T, Nakaya K, Lee VM, Trojanowski JQ, Iwatsubo T (1998) Aggregation of alpha-synuclein in Lewy bodies of sporadic Parkinson's disease and dementia with Lewy bodies. *Am J Pathol* 152(4):879–884
- Fujiwara H, Hasegawa M, Dohmae N, Kawashima A, Masliah E, Goldberg MS, Shen J, Takio K, Iwatsubo T (2002) Alpha-Synuclein is phosphorylated in synucleinopathy lesions. *Nat Cell Biol* 4(2):160–164, doi:10.1038/ncb748
- Polymeropoulos MH, Lavedan C, Leroy E, Ide SE, Dehejia A, Dutra A, Pike B, Root H, Rubenstein J, Boyer R, Stenroos ES, Chandrasekharappa S, Athanassiadou A, Papapetropoulos T, Johnson WG, Lazzarini AM, Duvoisin RC, Di Iorio G, Golbe LI, Nussbaum RL (1997) Mutation in the alpha-synuclein gene identified in families with Parkinson's disease. *Science* 276(5321):2045–2047
- Kruger R, Kuhn W, Muller T, Woitalla D, Graeber M, Kosel S, Przuntek H, Epplen JT, Schols L, Riess O (1998) Ala30Pro mutation in the gene encoding alpha-synuclein in Parkinson's disease. *Nat Genet* 18(2):106–108, doi:10.1038/ng0298-106
- Zarranz JJ, Alegre J, Gomez-Esteban JC, Lezcano E, Ros R, Ampuero J, Vidal L, Hoenicka J, Rodriguez O, Atares B, Llorens V, Gomez Tortosa E, Del Ser T, Munoz DG, de Yébenes JG (2004) The new mutation, E46K, of alpha-synuclein causes Parkinson and Lewy body dementia. *Ann Neurol* 55(2):164–173, doi:10.1002/ana.10795
- Appel-Cresswell S, Vilarino-Guelli C, Encarnacion M, Sherman H, Yu J, Shah B, Weir D, Thompson C, Szu-Tu C, Trinh J, Aasly JO, Rajput A, Rajput AH, Jon Stoessl A, Farrer MJ (2013) Alpha-synuclein p.H50Q, a novel pathogenic mutation for Parkinson's disease. *Mov Disord* 28(6):811–813, doi:10.1002/mds.25421
- Lesage S, Anheim M, Letournel F, Bousset L, Honore A, Rozas N, Pieri L, Madiona K, Durr A, Melki R, Verny C, Brice A (2013) G51D alpha-synuclein mutation causes a novel parkinsonian-pyramidal syndrome. *Ann Neurol* 73(4):459–471, doi:10.1002/ana.23894
- Singleton AB, Farrer M, Johnson J, Singleton A, Hague S, Kachergus J, Hulihan M, Peuralinna T, Dutra A, Nussbaum R, Lincoln S, Crawley A, Hanson M, Maraganore D, Adler C, Cookson MR, Muentner M, Baptista M, Miller D, Blacato J, Hardy J, Gwinn-Hardy K (2003) Alpha-Synuclein locus triplication causes Parkinson's disease. *Science* 302(5646):841, doi:10.1126/science.1090278
- Chartier-Harlin MC, Kachergus J, Roumier C, Mouroux V, Douay X, Lincoln S, Leveque C, Larvor L, Andrieux J, Hulihan M, Waucquier N, Defebvre L, Amouyel P, Farrer M, Destee A (2004) Alpha-synuclein locus duplication as a cause of familial Parkinson's disease. *Lancet* 364(9440):1167–1169, doi:10.1016/S0140-6736(04)17103-1
- Ibanez P, Bonnet AM, Debarges B, Lohmann E, Tison F, Pollak P, Agid Y, Durr A, Brice A (2004) Causal relation between alpha-synuclein gene duplication and familial Parkinson's disease. *Lancet* 364(9440):1169–1171, doi:10.1016/S0140-6736(04)17104-3
- Conway KA, Harper JD, Lansbury PT (1998) Accelerated in vitro fibril formation by a mutant alpha-synuclein linked to early-onset Parkinson disease. *Nat Med* 4(11):1318–1320, doi:10.1038/3311
- Choi W, Zibae S, Jakes R, Serpell LC, Davletov B, Crowther RA, Goedert M (2004) Mutation E46K increases phospholipid binding and assembly into filaments of human alpha-synuclein. *FEBS Lett* 576(3):363–368, doi:10.1016/j.febslet.2004.09.038
- Ghosh D, Mondal M, Mohite GM, Singh PK, Ranjan P, Anoop A, Ghosh S, Jha NN, Kumar A, Maji SK (2013) The Parkinson's disease-associated H50Q mutation accelerates alpha-Synuclein aggregation in vitro. *Biochemistry* 52(40):6925–6927, doi:10.1021/bi400999d
- Yonetani M, Nonaka T, Masuda M, Inukai Y, Oikawa T, Hisanaga S, Hasegawa M (2009) Conversion of wild-type alpha-synuclein into mutant-type fibrils and its propagation in the presence of A30P mutant. *J Biol Chem* 284(12):7940–7950, doi:10.1074/jbc.M807482200
- Braak H, Del Tredici K, Rub U, de Vos RA, Jansen Steur EN, Braak E (2003) Staging of brain pathology related to sporadic Parkinson's disease. *Neurobiol Aging* 24(2):197–211
- Attems J, Walker L, Jellinger KA (2014) Olfactory bulb involvement in neurodegenerative diseases. *Acta Neuropathol* 127(4):459–475, doi:10.1007/s00401-014-1261-7
- Luk KC, Kehm V, Carroll J, Zhang B, O'Brien P, Trojanowski JQ, Lee VM (2012) Pathological alpha-synuclein transmission initiates Parkinson-like neurodegeneration in nontransgenic mice. *Science* 338(6109):949–953, doi:10.1126/science.1227157
- Masuda-Suzukake M, Nonaka T, Hosokawa M, Oikawa T, Arai T, Akiyama H, Mann DM, Hasegawa M (2013) Prion-like spreading of pathological alpha-synuclein in brain. *Brain* 136(Pt 4):1128–1138, doi:10.1093/brain/awt037
- Mougenot AL, Nicot S, Bencsik A, Morignat E, Verchere J, Lakhdar L, Legastelois S, Baron T (2011) Prion-like acceleration of a synucleinopathy in a transgenic mouse model. *Neurobiol Aging*. doi:10.1016/j.neurobiolaging.2011.06.022
- Luk KC, Kehm VM, Zhang B, O'Brien P, Trojanowski JQ, Lee VM (2012) Intracerebral inoculation of pathological alpha-synuclein initiates a rapidly progressive neurodegenerative alpha-synucleinopathy in mice. *J Exp Med* 209(5):975–986, doi:10.1084/jem.20112457
- Watts JC, Giles K, Oehler A, Middleton L, Dexter DT, Gentleman SM, DeArmond SJ, Prusiner SB (2013) Transmission of multiple system atrophy prions to transgenic mice. *Proc Natl Acad Sci U S A* 110(48):19555–19560, doi:10.1073/pnas.1318268110
- Recasens A, Dehay B, Bove J, Carballo-Carbajal I, Dovero S, Perez-Villalba A, Fernagut PO, Blesa J, Parent A, Perier C, Farinas I, Obeso JA, Bezard E, Vila M (2014) Lewy body extracts from Parkinson disease brains trigger alpha-synuclein pathology and neurodegeneration in mice and monkeys. *Ann Neurol* 75(3):351–362, doi:10.1002/ana.24066
- Obi K, Akiyama H, Kondo H, Shimomura Y, Hasegawa M, Iwatsubo T, Mizuno Y, Mochizuki H (2008) Relationship of phosphorylated alpha-synuclein and tau accumulation to Abeta deposition in the cerebral cortex of dementia with Lewy bodies. *Exp Neurol* 210(2):409–420, doi:10.1016/j.expneurol.2007.11.019
- Jakes R, Crowther RA, Lee VM, Trojanowski JQ, Iwatsubo T, Goedert M (1999) Epitope mapping of LB509, a monoclonal antibody directed against human alpha-synuclein. *Neurosci Lett* 269(1):13–16
- Hasegawa M, Arai T, Nonaka T, Kametani F, Yoshida M, Hashizume Y, Beach TG, Buratti E, Baralle F, Morita M, Nakano I, Oda T, Tsuchiya K, Akiyama H (2008) Phosphorylated TDP-43 in frontotemporal lobar degeneration and amyotrophic lateral sclerosis. *Ann Neurol* 64(1):60–70, doi:10.1002/ana.21425
- Abeliovich A, Schmitz Y, Farinas I, Choi-Lundberg D, Ho WH, Castillo PE, Shinsky N, Verdugo JM, Armanini M, Ryan A, Hynes M, Phillips H, Sulzer D, Rosenthal A (2000) Mice lacking alpha-synuclein display functional deficits in the nigrostriatal dopamine system. *Neuron* 25(1):239–252
- Lee HJ, Gallagher M, Holland PC (2010) The central amygdala projection to the substantia nigra reflects prediction error information in appetitive conditioning. *Learn Mem* 17(10):531–538, doi:10.1101/lm.1889510
- Smith AD, Bolam JP (1990) The neural network of the basal ganglia as revealed by the study of synaptic connections of identified neurones. *Trends Neurosci* 13(7):259–265
- Gibb WR (1997) Functional neuropathology in Parkinson's disease. *Eur Neurol* 38(Suppl 2):21–25
- Freundt EC, Maynard N, Clancy EK, Roy S, Bousset L, Sourigues Y, Covert M, Melki R, Kirkegaard K, Brahm M (2012) Neuron-to-neuron transmission of alpha-synuclein fibrils through axonal transport. *Ann Neurol* 72(4):517–524, doi:10.1002/ana.23747
- Rey NL, Petit GH, Bousset L, Melki R, Brundin P (2013) Transfer of human alpha-synuclein from the olfactory bulb to interconnected brain regions in mice. *Acta Neuropathol* 126(4):555–573, doi:10.1007/s00401-013-1160-3
- Ulusoy A, Rusconi R, Perez-Reueta BI, Musgrove RE, Helwig M, Winzen-Reichert B, Di Monte DA (2013) Caudo-rostral brain spreading of alpha-synuclein through vagal connections. *EMBO Mol Med* 5(7):1051–1059, doi:10.1002/emmm.201302475
- Braak H, Rub U, Gai WP, Del Tredici K (2003) Idiopathic Parkinson's disease: possible routes by which vulnerable neuronal types may be subject to neuroinvasion by an unknown pathogen. *J Neural Transm* 110(5):517–536, doi:10.1007/s00702-002-0808-2
- Liu L, Drouot V, Wu JW, Witter MP, Small SA, Clelland C, Duff K (2012) Trans-synaptic spread of tau pathology in vivo. *PLoS One* 7(2):e31302, doi:10.1371/journal.pone.0031302
- Iba M, Guo JL, McBride JD, Zhang B, Trojanowski JQ, Lee VM (2013) Synthetic tau fibrils mediate transmission of neurofibrillary tangles in a

- transgenic mouse model of Alzheimer's-like tauopathy. *J Neurosci* 33(3):1024–1037, doi:10.1523/JNEUROSCI.2642-12.2013
39. Iseki E, Togo T, Suzuki K, Katsuse O, Marui W, de Silva R, Lees A, Yamamoto T, Kosaka K (2003) Dementia with Lewy bodies from the perspective of tauopathy. *Acta Neuropathol* 105(3):265–270, doi:10.1007/s00401-002-0644-3
 40. Colom-Cadena M, Gelpi E, Charif S, Belbin O, Blesa R, Marti MJ, Clarimon J, Lleo A (2013) Confluence of alpha-synuclein, tau, and beta-amyloid pathologies in dementia with Lewy bodies. *J Neuropathol Exp Neurol* 72(12):1203–1212, doi:10.1097/NEN.0000000000000018
 41. Clinton LK, Blurton-Jones M, Myczek K, Trojanowski JQ, LaFerla FM (2010) Synergistic Interactions between Abeta, tau, and alpha-synuclein: acceleration of neuropathology and cognitive decline. *J Neurosci* 30(21):7281–7289, doi:10.1523/JNEUROSCI.0490-10.2010
 42. Guo JL, Covell DJ, Daniels JP, Iba M, Stieber A, Zhang B, Riddle DM, Kwong LK, Xu Y, Trojanowski JQ, Lee VM (2013) Distinct alpha-synuclein strains differentially promote tau inclusions in neurons. *Cell* 154(1):103–117, doi:10.1016/j.cell.2013.05.057
 43. Chung CY, Khurana V, Auluck PK, Tardiff DF, Mazzulli JR, Soldner F, Baru V, Lou Y, Freyzon Y, Cho S, Mungenast AE, Muffat J, Mitalipova M, Pluth MD, Jui NT, Schule B, Lippard SJ, Tsai LH, Krainc D, Buchwald SL, Jaenisch R, Lindquist S (2013) Identification and rescue of alpha-synuclein toxicity in Parkinson patient-derived neurons. *Science* 342(6161):983–987, doi:10.1126/science.1245296
 44. Volpicelli-Daley LA, Luk KC, Patel TP, Tanik SA, Riddle DM, Stieber A, Meaney DF, Trojanowski JQ, Lee VM (2011) Exogenous alpha-synuclein fibrils induce Lewy body pathology leading to synaptic dysfunction and neuron death. *Neuron* 72(1):57–71, doi:10.1016/j.neuron.2011.08.033
 45. Nonaka T, Watanabe ST, Iwatsubo T, Hasegawa M (2010) Seeded aggregation and toxicity of {alpha}-synuclein and tau: cellular models of neurodegenerative diseases. *J Biol Chem* 285(45):34885–34898, doi:10.1074/jbc.M110.148460

doi:10.1186/s40478-014-0088-8

Cite this article as: Masuda-Suzukake *et al.*: Pathological alpha-synuclein propagates through neural networks. *Acta Neuropathologica Communications* 2014 **2**:88.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

